BETAINE

HYDROCHLORIC ACID SOURCE

DIGESTIVE SUPPORT

650 MG • 180 TABLETS

Supplement Facts Serving Size 1 Tablet

	Amount Per Serving
taine Hydrochloride	650 mg†
nsin 1 · 3000	165 mot

†Daily Value not established.

Other ingredients: stearic acid, microcrystalline cellulose, hydroxypropyl cellulose, modified cellulose gum, and colloidal silicon dioxide.

Contains milk.

Suggested Use: 1 tablet with a meal, up to four times daily. Do not take on an empty stomach.

CAUTION: Do not use if you are pregnant, may become pregnant, or breastfeeding. If a burning sensation is experienced, discontinue use, or use less frequently.

WARNING: If you have stomach or duodenal ulcers, or are taking drugs that may cause ulcers, do not take this product.

In the stomach, hydrochloric acid converts pepsinogen to pepsin, an enzyme that breaks down protein. Pepsin breaks down proteins into smaller, more easily absorbed substances. As people age, hydrochloric acid secretion may be reduced, which can result in decreased levels of pepsin. Without proper pepsin levels, the body has a difficult time digesting food. Betaine hydrochloride is used as a supplemental source of hydrochloric acid, which can be utilized by the stomach to produce pepsin. The combination of betaine hydrochloride and pepsin provide an excellent stomach tonic.*

Do not use if either tamper-evident seal is broken or missing. Keep out of the reach of children.

Contains no yeast, egg, gluten, corn, soy or wheat. Contains no sugar, starch, salt, preservatives, or artificial color, flavor or fragrance.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

SN1362 REV B112